Column name	Explanation
	(data type, levels)
	eline data
Gender	Gender (string, male/female)
age & age_collapsed	Patients' age in years, additionally grouped into 5
	year ranges (integer)
diagnosis & diagnosis_collapsed	Neurological main diagnosis (string), for ease of
	use additionally grouped into cerebrovascular
Marital_status & marital_status_collapsed	disorders, neuromuscular disorders, epilepsy,
	movement disorders and others (string)
	Marital state (string, married; single;
	widowed/divorced), additionally reduced to
	married or not married
Living_situation	Living situation (string, alone or not alone, e.g.
	with partner or family)
education	Education (string, high: German abitur or
	university, ≥ 12 years; medium: German
	Realschule or general certificate of secondary
	education, 8 to 12 years; and low: German
Addata and addition	Hauptschule or no school ≤ 8 years)
tablets_preparation	Preparation and taking of medication
	independently (string, independent vs needs help from others)
number of drugs per day	,
number of drugs per day	total amount of different medications taken daily
come 1 to com 10	(integer) Items for Stendal Adherence to Medication Score
sams_1 to sam_18	(SAMS), multinomial/integer, range 0 – 4, with
	higher numbers indicating higher levels of
	nonadherence
SAMS	Stendal Adherence to Medication Score (SAMS)
SAIVIS	total sum score, integer
bdi_1 to bdi_21	Items for Beck's Depression Inventory (BDI) II,
501_1 to 501_21	multinomial/integer, range 0 - 3, with higher
	numbers indicating higher intensity of depressive
	symptoms
BDI	Beck's Depression Inventory (BDI) II sum score,
	integer
	Note: values with * were taken from
	neuropsychological testing performed in routine
	assessment on the wards, not by study staff, hence
	only sum scores are available
bfi_1 to bfi_10	Items of Big Five Inventory (BFI) 10,
	multinomial/integer, range 1 – 5 with 1 = I do not
	agree at all and 5 = I completely agree
BFI	Personality according to Big Five Inventory (BFI) 10
	(string, agreeableness, conscientiousness,
	extraversion, openness, neuroticism)
hccq_1 to hccq_15	Items of Healthcare Climate Questionnaire (HCCQ),
	multinomial/integer, range 1 -7 with 1 = I do not
	agree at all to 7 = I completely agree
HCCQ	Sum Score of Health Care Climate Questionnaire
	(HCCQ) (numeric/float)

Moca_1 to moca_13f	Items for Montreal Cognitive Assessment (MoCA)
	1: visuospatial/executive, connecting numbers
	2: visuospatial/executive, drawing cube
	3: visuospatial/executive, clock a) contour b)
	numbers c) hands
	4: naming animals
	5: attention, repeating numbers forward
	6, attention, repeating numbers backwards7,
	7, attention, list of letters
	8, attention, calculation
	9, language, repeating sentence
	10, language, word naming
	11, abstraction
	12, delayed recall
	13, orientation a) day b) month c) year d) weekday
	e) location f) city
	Note: values with * were taken from
	neuropsychological testing performed in routine
	assessment on the wards, not by study staff, hence
	only sum scores are available
moca_edu_extra	Additional point for less than 12 years of education
	Sum Score for Montreal Cognitive Assessment,
MoCA	
TuC Cocondo and TuC	integer, range 19-30
TuG_Seconds and TuG	Timed Up and Go (TuG) duration in seconds,
	integer
	Additionally grouped into grouped into <20 sec, 20
	- 30 sec, > 30 sec, and inability to perform the test
	due to medical reasons (string)
doc_app_frequency_quarterly	Number of doctor consultations per quarter
	(float)
Walking Aid	Use of any form of walking aid, e.g. wheelchair,
	walking stick, crutches etc.
	(string, yes/no)
nonmedical_treat	Use of any nonmedical treatment, e.g.
	physiotherapy, occupational therapy, or speech
	therapy
	(string, yes/no)
frequency of consultation quarterly	Number of doctor consultations per quarter
	(float)
medication_change	Has there been a change in medication in the last 6
-	months?
	(string, yes/no)
medic_change_type	What kinds of changes were made? "Cannot
	provide information" = patient reports a change
	was made but cannot name what was changed
	(string, new medicament, reduction, increase,
	medicament discontinued, other)
Sf_1 to sf_11d	Items for Short-Form 36 (SF-36),
01 00 01_110	multinomial/integer, range 1 – 5 with 1 = I do not
	agree at all and 5 = I completely agree
rphysf_c	SF-36 domain Physical functioning
1611831_c 	(integer)
	(miteger)

rsocf_c	SF-36 domain Social functioning (float)
rrolef_c	SF-36 domain Role limitations due to physical health
	(integer)
	SF-36 domain Role limitations due to emotional
rrolee_c	
	problems
	(integer)
rment_c	SF-36 domain Emotional well-being
	(integer)
rvit_c	SF-36 domain Energy/fatigue
	(integer)
rpain_c	SF-36 domain Pain
	(float)
rgenh_c	SF-36 domain General health
	(integer)
rhchange_c	SF-36 domain Health change
	(integer)
SF_36_PCS	SF-36 Physical Component Scale
	(float)
SF_36_MCS	SF-36 Mental Component Scale
	(float)
Follow-Up interview 1 mo	nth after discharge from hospital
geradh4_FUI_medication_change	Follow-Up I Item, Change of medication since
Bernen T. e.T. enemen Terran Be	discharge from hospital
	(string, yes/no)
geradh6_FUI_medication_change_by_whom	Follow-Up I Item, Change of medication issued by
geraano_r or_meateation_change_by_whom	whom?
	(string, physician, patient without consulting
	physician, other)
geradh7_FUI_medication_change_reason	Follow-Up I Item, reason for change of medication
80.00	(string, new indication, no effect, side effects,
	polymedication (too many prescriptions), other)
Follow-Un interview 12 mo	nths after discharge from hospital
geradh4_FU2_medication_change	Follow-Up II Item, Change of medication in the last
geraun4_roz_medication_change	year?
	'
geradh6_FU2_medication_change_by_who	(string, yes/no) Follow-Up II Item, Change of medication issued by
	whom?
m	
	(string, physician, patient without consulting physician, other)
goradh7 EU2 modication change reases	
geradh7_FU2_medication_change_reason	Follow-Up II Item, reason for change of medication
	(string, new indication, no effect, side effects,
2000 4 511	polymedication (too many prescriptions), other)
sams_1_FU	Follow-Up II SAMS Item 2, Do you know the reason
	for taking your medication?
2.511	integer, range 0-4, 0 = for none, 4 = for all
sams_2_FU	Follow-Up II SAMS Item 2, Do you know the
	dosages of your medication?
	integer, range 0-4, 0 = for all, 4 = for none
sams_4_FU	integer, range 0-4, 0 = for all, 4 = for none Follow-Up II SAMS Item 4, Do you take your medication regularly?

	integer, range 0-4, 0 = all, 4 = none
sams_8_FU	Follow-Up II SAMS Item 8, Do you stop taking your
	medication when you feel better?
	integer, range 0-4, 0 = never, 4 = most of the time
sams_11_FU	Follow-Up II SAMS Item 11, If you think you have
	side effects due to of the medications (such as
	tremors, nausea etc.), do you reduce the dose
	without consulting a doctor?
	integer, range 0-4, 0 = never, 4 = most of the time
sams_12_FU	Follow-Up II SAMS Item 12, If you think you have
	side effects due to of the medications (such as
	tremors, nausea etc.), do you not take the
	medication for a while, i.e. take a break?
	integer, range 0-4, 0 = never, 4 = most of the time
sams_13_FU	Follow-Up II SAMS Item 13, If you feel you have to
	take too many, do you stop taking those
	medications you consider to be less important
	than the others without consulting your doctor?
	integer, range 0-4, 0 = never, 4 = most of the time
sams_14_FU	Follow-Up II SAMS Item 14, If you forget or omit
	your medication, do you forget it in the morning?
	integer, range 0-4, 0 = never, 4 = most of the time
sams_15_FU	Follow-Up II SAMS Item 15, If you forget or omit
	your medication, do you forget it at noon?
	integer, range 0-4, 0 = never, 4 = most of the time
sams_16_FU	Follow-Up II SAMS Item 16, If you forget or omit
	your medication, do you forget it in the evening?
	integer, range 0-4, 0 = never, 4 = most of the time
SF12_1	SF-12 general health
	(integer, 1 = excellent, 2 = very good, 3 = good, 4 =
	fair, 5 = poor)
SF12_2	SF-12 limited in moderate activities
	(integer, 1 = limited a lot, 2 = limited a little, 3 =
	not limited at all)
SF12_3	SF-12 limited in climbing several flights of stairs
	(integer, 1 = limited a lot, 2 = limited a little, 3 =
	not limited at all)
SF12_4	SF-12 accomplishing less (physical)
	(integer, 1 = yes, 2 = no)
SF12_5	SF-12 limited in the kind of work or other activities
	(physical)
5540.6	(integer,
SF12_6	SF-12 accomplishing less (emotional)
5542.7	(integer, 1 = yes, 2 = no)
SF12_7	SF-12 less carefully (emotional)
5542.0	(integer, 1 = yes, 2 = no)
SF12_8	SF-12 interference of pain with work
	(integer, 1 = not at all, 2 = a little bit, 3 =
	moderately, 4 = quite a bit, 5 = extremely)
SF12_9	SF-12 felt calm and peaceful

	(integer, 1 = all of the time, 2 = most of the time, 3 = a bit of the time, 4 = some of the time, 5 = a little bit of the time, 6 = none of the time)
SF12_10	SF-12 had lot of energy (integer, 1 = all of the time, 2 = most of the time, 3 = a bit of the time, 4 = some of the time, 5 = a little bit of the time, 6 = none of the time)
SF12_11	SF-12 felt downhearted and blue (integer, 1 = all of the time, 2 = most of the time, 3 = a bit of the time, 4 = some of the time, 5 = a little bit of the time, 6 = none of the time)
SF12_12	SF-12 time of interference with social activities integer, 1 = all of the time, 2 = most of the time, 3 = some of the time, 4 = a little bit of the time, 5 = none of the time)

References

- 1. Beck, A., Steer, R. & Brown, G. Beck depression inventory—second edition: manual. *San Antonio: The Psychological Corporation* **4**, 561-571 (1996).
- 2. Hautzinger, M., Keller, F. & Kühner, C. *Beck-Depressions-Inventar: Revision*. (Harcourt test services, 2006).
- 3. Nasreddine, Z. S. *et al.* The Montreal Cognitive Assessment, MoCA: a brief screening tool for mild cognitive impairment. *J Am Geriatr Soc* **53**, 695-699, doi:10.1111/j.1532-5415.2005.53221.x (2005).
- 4. Podsiadlo, D. & Richardson, S. The timed "Up & Go": a test of basic functional mobility for frail elderly persons. *J Am Geriatr Soc* **39**, 142-148, doi:10.1111/j.1532-5415.1991.tb01616.x (1991).
- 5. John, O. P., Donahue, E. M. & Kentle, R. L. *The Big Five Inventory--Versions 4a and 54*. (University of California, Berkeley, Institute of Personality and Social Research, 1991).
- Rammstedt, B. The 10-item Big Five Inventory: Norm values and investigation of sociodemographic effects based on a German population representative sample. European Journal of Psychological Assessment 23, 193-201, doi:10.1027/1015-5759.23.3.193 (2007).
- 7. Schmidt, K. *et al.* Autonomy support in primary care—validation of the German version of the Health Care Climate Questionnaire. *Journal of Clinical Epidemiology* **65**, 206-211, doi:10.1016/j.jclinepi.2011.06.003 (2012).
- 8. Gremigni, P. Validation of the Health Care Communication Questionnaire (HCCQ) to measure patients evaluation of hospital personnel communication skills. *Patient Education and Counseling* **71**, 57-64 (2008).
- 9. Ware, J. E., Jr. & Sherbourne, C. D. The MOS 36-item short-form health survey (SF-36). I. Conceptual framework and item selection. *Med Care* **30**, 473-483 (1992).
- Ware, J., Jr., Kosinski, M. & Keller, S. D. A 12-Item Short-Form Health Survey: construction of scales and preliminary tests of reliability and validity. *Med Care* 34, 220-233, doi:10.1097/00005650-199603000-00003 (1996).
- 11. Franke, G. H., Nentzl, J. & Jagla-Franke, M. SAMS. Stendal Adherence to Medication Score. Testmanual. www.psychometrikon.de, doi:10.6099/1000372 (2020).